

AD

\_\_\_\_\_  
(Leave blank)

Award Number: W81XWH-08-1-0763

TITLE: Characterization of IKBKE as a Breast Cancer Oncogene

PRINCIPAL INVESTIGATOR: Alicia Zhou

CONTRACTING ORGANIZATION: Dana-Farber Cancer Institute  
Boston, MA 02115

REPORT DATE: October 2009

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

☒ Approved for public release; distribution unlimited

☐ Distribution limited to U.S. Government agencies only;  
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 10/31/2009		2. REPORT TYPE Annual Summary		3. DATES COVERED (From - To) 10/01/2008 - 09/30/2009	
4. TITLE AND SUBTITLE Characterization of IKBKE as a Breast Cancer Oncogene			5a. CONTRACT NUMBER W81XWH-08-1-0763		
			5b. GRANT NUMBER BC083046-PreDoc		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Alicia Zhou Email: alicia_zhou@dfci.harvard.edu			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Dana-Farber Cancer Institute Boston, MA 02115			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research And Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Previous work in the Hahn Lab has identified <i>IKBKE</i> as a novel breast cancer oncogene that is capable of human mammary cell transformation. Amplification and overexpression of IKKepsilon was observed in a significant percentage of human breast cancer cell lines and primary tumor samples. Additionally, breast cancer cell lines carrying the <i>IKBKE</i> amplicon showed decreased viability in response to IKKepsilon suppression by shRNA. Our work has also demonstrated that IKKepsilon is a non-canonical IKK (IkappaB kinase) family member that activates the NF-kappaB pathway and that this activity is essential for cell transformation. However, it is known that IKKepsilon does not participate in the canonical IKK complex to activate NF-kappaB signaling. Though recent work has sought to identify the downstream targets of IKKepsilon, the mechanism of its upstream regulation is not well-understood. Thus, I propose to further our understanding of IKKepsilon function by investigating the upstream regulation of IKKepsilon – specifically, the role of ubiquitination in IKKepsilon-mediated cell transformation. In addition, I am to investigate the role of IKKepsilon in breast cancer initiation and maintenance <i>in vivo</i> through the generation of a constitutive and an inducible transgenic mouse model.					
15. SUBJECT TERMS breast cancer, ubiquitin, transgenic mouse model, IKKepsilon, NF-kappaB					
16. SECURITY CLASSIFICATION OF: U			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  8	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusion.....	7

## Introduction


I have made significant progress in both Specific Aims that were originally laid out in my Statement of Work. In this Progress Report, I will report the studies and results that I have conducted over the past year towards accomplishing those aims.

## Specific Aims

1. Investigate the role of ubiquitination in IKK $\epsilon$ -mediated cell transformation
  - a. Confirm and characterize IKK $\epsilon$  ubiquitination in the context of mammary cell transformation
  - b. Identification of the IKK $\epsilon$  ubiquitin-accepting residues
  - c. Determine the functional relevance of IKK $\epsilon$  ubiquitination in mammary cell transformation
2. Investigate the role of IKK $\epsilon$  in breast cancer initiation and maintenance
  - a. Investigate the role of *IKBKE* in breast cancer initiation
  - b. Investigate the role of *IKBKE* in breast cancer maintenance


## Studies and Results

**Specific Aim 1a:** I have confirmed that IKK $\epsilon$  ubiquitination occurs in a mammary epithelial cell context. I immunoprecipitated IKK $\epsilon$  from human mammary epithelial cells that stably overexpress activated MEK (HMECM) along with either Flag-tagged (F-IKK $\epsilon$ ) or myristolated Flag-tagged (MF-IKK $\epsilon$ ) IKK $\epsilon$  and was able to detect an endogenously ubiquitinated species of MF-IKK $\epsilon$  by immunoblotting for ubiquitin (Figure 1A). In parallel, I transiently transfected the same cells with HA-tagged ubiquitin, performed an immunoprecipitation for IKK $\epsilon$  and was able to successfully detect an ubiquitinated ladder of MF-IKK $\epsilon$  by immunoblotting for HA. I performed the reverse IP/Western experiment, and was also able to immunoblot for IKK $\epsilon$  after immunoprecipitation by HA antibody (Figure 1B). Interestingly, I was only able to confirm IKK $\epsilon$  ubiquitination in the MF-IKK $\epsilon$  transformed HMECM cells and not in the F-IKK $\epsilon$  transformed cells. Previous characterization of these cell lines has shown that MF-IKK $\epsilon$  exhibits a much more robust transformation phenotype in HMECM cells than the F-IKK $\epsilon$  counterpart. Perhaps this variation in transformation phenotype is related to the ubiquitination status of IKK $\epsilon$  in these cells.




I have attempted to characterize the lysine linkage of IKKε using K48- or K63- only ubiquitin mutants. However, I have thus far found that these ubiquitin mutants have yielded inconclusive immunoprecipitation and immunoblot results. This has made it difficult to specifically characterize the nature of the interaction. I am currently working on optimizing the conditions for the use of these ubiquitin mutant constructs.

**Specific Aim 1b:** I have successfully identified three lysine residues in IKKε that are subject to ubiquitination by mass spectrometry. I transiently cotransfected GST-tagged IKKε and HA-tagged ubiquitin in HEK293T cells and performed a GST immunoprecipitation. The immunoprecipitates were then



subjected to SDS-PAGE followed by Coomassie blue staining. Four bands of interest were identified and submitted for mass spectrometry analysis (Figure 2). We obtained 58.2% coverage of the IKKε protein and 64.7% (22 out of 34) coverage of the internal lysines. From this analysis, three lysine residues were identified as modified by ubiquitin: K30, K401, and K416.


**Specific Aim 1c:** I have successfully generated site-specific lysine-to-alanine IKKε mutants for the three lysine residues that were identified in Aim 1b. All possible combinations of single, double and triple mutants of the protein were generated and then retrovirally introduced into HA1EM cell to create stable cell lines that express these mutant constructs. These mutants were assessed for transformation capacity by soft agar analysis (Figure 3). From this analysis, I have determined that the K30 and K401 residues are essential for IKKε-mediated transformation. However mutation of the K416 residue does not perturb the transformation phenotype of IKKε. Surprisingly, in addition, I have observed that the mutation of the K416 residue rescues the decreased transformation phenotype of the K30 and K401 mutants. This may be due to a



combination of positive and negative regulation of IKK $\epsilon$  by both Lysine48- and Lysine63- linked ubiquitination.


In order to assess the tumorigenicity capacity of these mutants, all mutant IKK $\epsilon$  cell lines have been injected into immunocompromised mice (6 injections per cell line) and are currently being observed for tumor formation.

**Specific Aim 2a:** I have successfully generated a MMTV-IKK $\epsilon$  transgenic mouse model. By genotyping analysis, 9 founder mice were identified. These founders were bred out and assessed for mammary gland-specific expression of IKK $\epsilon$  by immunoblot (Figure 4). From this analysis, 3 founder lines with varying levels of IKK $\epsilon$  expression were chosen and are currently being expanded and followed. The original founders are now ~8 months old and have not shown a mammary tumor formation phenotype.



In addition, these founder lines are also being bred to MMTV-ErbB2 mice, which are known to have a delayed mammary gland tumor formation phenotype. These MMTV-IKK $\epsilon$ /ErbB2 bitransgenic mice will allow us to determine if there is a synergistic tumor formation phenotype, should the MMTV-IKK $\epsilon$  mice not develop tumors on their own. We have also obtained WAP-Cre and p53 floxed mice from the Mouse Models of Human Cancer Consortium (MMHCC) and are prepared to cross the MMTV-IKK $\epsilon$  mice into a mammary-specific p53 null background, if needed.

**Specific Aim 2b:** We have successfully generated a doxycycline-inducible MMTV-rtTA/TetO-IKK $\epsilon$  bitransgenic mouse model. By genotyping analysis, 5 founder mice were identified (Figure 5). After analysis for doxycycline-inducible mammary gland IKK $\epsilon$  expression, 2 founder lines with varying levels of IKK $\epsilon$  expression were chosen and are currently being expanded and followed.



## **Key Research Accomplishments**

- Confirmed IKK $\epsilon$  ubiquitination in the context of mammary cell transformation
- Identified three lysine residues within IKK $\epsilon$  that are subject to ubiquitination by mass spectrometry
- Generated IKK $\epsilon$  point mutants in which the identified lysine residues are mutated singularly and in all possible tandem combinations
- Generated cell lines that stably express the IKK $\epsilon$  mutant constructs
- Demonstrated that IKK $\epsilon$  lysine mutants show a distinct change in transformation phenotype *in vitro*
- Generated founder mice for the constitutive MMTV-IKK $\epsilon$  transgenic mouse model
- Generated founder mice for the inducible MMTV-rtTA/TetO-IKK $\epsilon$  transgenic mouse model

## **Reportable Outcomes**

- Developed 14 HA1EM mutant IKK $\epsilon$  cell lines: both F-IKK $\epsilon$  and MF-IKK $\epsilon$  with the following mutations K30A; K401A; K416A; K30A K401A; K30A K416A; K401A K416A; K30A K401A K416A
- Developed two new animal models: MMTV-IKK $\epsilon$  and MMTV-rtTA/TetO-IKK $\epsilon$

## **Conclusion**

Over the past year, I have made significant progress in all aspects of my Specific Aims as laid out in my original Statement of Work.

I have been able to identify three residues of IKK $\epsilon$  that undergo modification by ubiquitination, and I have been able to show that the mutation of these residues is able to perturb the IKK $\epsilon$ -mediated transformation phenotype. However, my mutational studies have also indicated that there is a more complex and dynamic role for IKK $\epsilon$  ubiquitination than I had originally hypothesized. I believe that IKK $\epsilon$  is undergoing a combination of both Lys48- and Lys63- linked ubiquitination, and that this combination of modifications is serving to tightly regulate both the levels of IKK $\epsilon$  protein in the cell as well as its enzymatic activation. I propose to do a series of IP/Western experiments that utilize the Lys48- and Lys63- only ubiquitin mutants in combination with the IKK $\epsilon$  mutants, in order to determine the exact nature of these modifications. In addition, to understand how the perturbation of the various residues of IKK $\epsilon$  affects its enzymatic activity, I propose to perform *in vitro* kinase assays to assess the enzymatic activity of the various IKK $\epsilon$  mutants. Finally, I will also assay for NF- $\kappa$ B activation in these mutant cells to determine how the perturbation of IKK $\epsilon$  ubiquitination affects its role in this pathway.

I have also successfully generated both a constitutive and inducible transgenic mouse model of mammary-specific IKK $\epsilon$  expression. As originally laid out in my Statement of Work, I am currently working to generate a cohort of 50 mice to assess for a mammary tumor formation phenotype. In parallel, I have bred the constitutive MMTV-IKK $\epsilon$  with MMTV-ErbB2 mice to determine if there is a synergistic tumor formation phenotype, should the MMTV-IKK $\epsilon$  mice not develop

tumors on their own. In addition, we also have WAP-Cre and p53 floxed mice and are prepared to cross the MMTV-IKK $\epsilon$  mice into a mammary-specific p53 null background, if needed.